arXiv:1404.6982v1 [math.RT] 28 Apr 2014

Abstract Harmonic Analysis on the General Affine Group $GA(n,\mathbb{R})$

Kahar El-Hussein
Department of Mathematics, Faculty of Science,
Al Furat University, Dear El Zore, Syria and
Department of Mathematics, Faculty of Arts Science Al Quryyat,
Al-Jouf University, KSA
E-mail : kumath@ju.edu.sa, kumath@hotmail.com

January 16, 2016

Abstract

Let $GL(n,\mathbb{R})$ be the general linear group and let $GA(n,\mathbb{R}) = \mathbb{R}^n \rtimes_\rho GL(n,\mathbb{R})$ be its general affine group. Let $GL_+(n,\mathbb{R})$ be the identity component of $GL(n,\mathbb{R})$, which consists of the real $n\times n$ matrices with positive determinant and let $GL_-(n,\mathbb{R})$ be the set of all matrices with negative determinant. Since $GL(n,\mathbb{R})$ is a two copies of $GL_+(n,\mathbb{R})$, *i.e* $GL(n,\mathbb{R}) = GL_+(n,\mathbb{R}) \cup GL_+(n,\mathbb{R})$, because $GL_-(n,\mathbb{R})$ has a structure of group isomorphic onto $GL_+(n,\mathbb{R})$, see [13]. Therefore first we consider the group $GA_+(n,\mathbb{R}) = \mathbb{R}^n \rtimes_\rho GL_+(n,\mathbb{R})$ to generalize the Fourier transform and to prove the Plancherel theorem for $GA_+(n,\mathbb{R})$. Secondly and since $GA(n,\mathbb{R})$ is a two copies of $GA_+(n,\mathbb{R})$, so we can easily to establish the Plancherel theorem for $GA(n,\mathbb{R})$.

Keywords: Linear Group $GL(n,\mathbb{R})$, Affine General Group $GA(n,\mathbb{R})$, Fourier Transform, Plancherel Theorem
AMS 2000 Subject Classification: $43A30\&35D$ 05

1 Introduction.

1.1. The general affine group $GA(n,\mathbb{R}) = \mathbb{R}^n \rtimes_\rho GL(n,\mathbb{R})$ of the general linear group $GL(n,\mathbb{R})$, which is the semidirect of the real vector $\mathbb{R}^n$ with $GL(n,\mathbb{R})$. Another kind of the affine group is the Poincare group $\mathbb{R}^4 \rtimes O(3.1)$, which is the affine group of the inhomogeneous Lorentz group $O(3.1)$. As well known the affine groups play an important role in physics in cosmology, gauge theory, gravitation, general relativity, ect... .The spacetime symmetry of the affine modeless of gravity is given by the general affine group $GA(n,\mathbb{R})$. The geometrical arena of the theory gravitation and electromagnetism is the modified affine frame bundle over a four dimensional spacetime manifold M, the structure group of the frame bundle is the affine group $GA(4,\mathbb{R}) = \mathbb{R}^4 \rtimes GL(4,\mathbb{R})$. The usual definition of phase space coordinates in terms of linear frames and use affine frames instead. This leads from $GL(4,\mathbb{R})$ covariance to $GA(4,\mathbb{R})$ covariance and means that the bundle of linear frames, is replaced by the bundle of affine frames. Also the affine group $GA(2,\mathbb{R})$ has a fundamental role in the visualization. One asks can we do the non commutative Fourier analysis on the group $GA(n,\mathbb{R}) = \mathbb{R}^n \rtimes GL(n,\mathbb{R})$. In fact that the abstract harmonic analysis on the locally compact groups is generally a difficult task. Still now neither the theory of quantum groups nor the representations theory have done to reach this goal. Recently, these problems found a satisfactory solution with the papers [9, 10, 11]. In this paper we will generalize our methods in [12, 13] to define the Fourier transform and to establish the Plancherel theorem for the general affine group on $GA(n,\mathbb{R})$.

2 Notation and Results for the Nilpotent Lie Group N.

2.1. The fine structure of the nilpotent Lie groups will help us to do the Fourier transform on a connected and simply connected nilpotent Lie groups N. As well known any group connected and simply connected N has the

following form

$$N = \begin{pmatrix} 1 & x_1^1 & x_1^2 & x_1^3 & . & . & . & . & . & x_1^{n-2} & x_1^{n-1} & x_1^n \\ 0 & 1 & x_2^2 & x_2^3 & . & . & . & . & . & x_2^{n-2} & x_2^{n-1} & x_2^n \\ 0 & 0 & 1 & x_3^3 & . & . & . & . & . & x_3^{n-2} & x_3^{n-1} & x_3^n \\ 0 & 0 & 0 & 1 & . & . & . & . & . & x_4^{n-2} & x_4^{n-1} & x_4^n \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & x_{n-2}^{n-2}. & .x_{n-2}^{n-1} & x_{n-2}^n \\ 0 & 0 & 0 & 0 & . & . & . & . & . & 1 & x_{n-1}^{n-2} & x_{n=1}^n \\ 0 & 0 & 0 & 0 & . & . & . & . & . & 0 & 1 & x_n^n \\ 0 & 0 & 0 & 0 & . & . & . & . & . & 0 & 0 & 1 \end{pmatrix} \tag{1}$$

As shown, this matrix is formed by the subgroup $\mathbb{R}$, $\mathbb{R}^2$,...., $\mathbb{R}^{n-1}$, and $\mathbb{R}^n$

$$\begin{pmatrix} \mathbb{R} = \begin{bmatrix} x_1^1 \\ 1 \\ 0 \\ 0 \\ . \\ . \\ . \\ . \\ . \\ 0 \\ 0 \\ 0 \end{bmatrix}, \mathbb{R}^2 = \begin{bmatrix} x_1^2 \\ x_2^2 \\ 1 \\ 0 \\ . \\ . \\ . \\ . \\ . \\ 0 \\ 0 \\ 0 \end{bmatrix}, .., \mathbb{R}^{n-1} = \begin{bmatrix} x_1^{n-1} \\ x_2^{n-1} \\ x_3^{n-1} \\ x_4^{n-1} \\ . \\ . \\ . \\ . \\ .x_{n-2}^{n-1} \\ x_{n-1}^{n-2} \\ 1 \\ 0 \end{bmatrix}, \mathbb{R}^n = \begin{bmatrix} x_1^n \\ x_2^n \\ x_3^n \\ x_4^n \\ . \\ . \\ . \\ . \\ x_{n-2}^n \\ x_{n=1}^n \\ x_n^n \\ 1 \end{bmatrix} \end{pmatrix} \tag{2}$$

Each $\mathbb{R}^i$ is a subgroup of N of dimension i , $1 \leq i \leq n$, put $d = n + (n-1) + ... + 2 + 1 = \frac{n(n+1)}{2}$, which is the dimension of N . According to [9], the group N is isomorphic onto the following group

$$(((((\mathbb{R}^n \rtimes_{\rho_n}) \mathbb{R}^{n-1}) \rtimes_{\rho_{n-1}}) \mathbb{R}^{n-2} \rtimes_{\rho_{n-2}} .....) \rtimes_{\rho_2} \mathbb{R}^2) \rtimes_{\rho_1} \mathbb{R} \tag{3}$$

That means

$$N \simeq (((((\mathbb{R}^n \rtimes_{\rho_n})\mathbb{R}^{n-1})\rtimes_{\rho_{n-1}})\mathbb{R}^{n-2} \rtimes_{\rho_{n-2}} .....) \rtimes_{\rho_4} \mathbb{R}^3 \rtimes_{\rho_3} \mathbb{R}^2) \rtimes_{\rho_2} \mathbb{R} \quad (4)$$

2.2. Denote by $L^1(N)$ the Banach algebra of N that consists of all complex valued functions on the group N, which are integrable with respect to the Haar measure of N and multiplication is defined by convolution on N as follows:

$$g * f(X) = \int_N f(Y^{-1}X)g(Y)dY \quad (5)$$

for any $f \in L^1(N)$ and $g \in L^1(N)$, where $X = (X^1, X^2, X^3, ..., X^{n-2}, X^{n-1}, X^n) \in N$, $Y = (Y^1, Y^2, Y^3, ..., Y^{n-2}, Y^{n-1}, Y^n) \in N$, $X^1 = x_1^1$, $X^2 = (x_1^2, x_2^2)$, $X^3 = (x_1^3, x_2^3, x_3^3)$, ..., $X^{n-2} = (x_1^{n-2}, x_2^{n-2}, x_3^{n-2}, x_4^{n-2}, ..., x_{n-2}^{n-2})$, $X^{n-1} = (x_1^{n-1}, x_2^{n-1}, x_3^{n-1}, x_4^{n-1}, ..., x_{n-2}^{n-1}, x_{n-1}^{n-1})$, $X^n = (x_1^n, x_2^n, x_3^n, x_4^n, ..., x_{n-2}^n, x_{n-1}^n, x_n^n)$ and $dY = dY^1dY^2dY^3, ..., dY^{n-2}dY^{n-1}dY^n$ is the Haar measure on N and $*$ denotes the convolution product on N. We denote by $L^2(N)$ its Hilbert space. We refer to [12] to define the Fourier transform on N

Definition 2.1. *For $1 \leq i \leq n$, let $\mathcal{F}^i$ be the Fourier transform on $\mathbb{R}^i$ and $0 \leq j \leq n-1$, let $\prod_{0\leq l\leq j} \mathbb{R}^{n-l} = (..((((\mathbb{R}^n \rtimes_{\rho_n})\mathbb{R}^{n-1})\rtimes_{\rho_{n-1}})\mathbb{R}^{n-2}\rtimes_{\rho_{n-2}})..\times_{\rho_{n-j}} \mathbb{R}^{n-j})$, and let $\prod_{0\leq l\leq j} \mathcal{F}^{n-l} = \mathcal{F}^n\mathcal{F}^{n-l}\mathcal{F}^{n-2}....\mathcal{F}^{n-j}$, we can define the Fourier transform on $\prod_{0\leq l\leq n-1} \mathbb{R}^{n-l} = \mathbb{R}^n \rtimes_{\rho_n} \mathbb{R}^{n-1} \rtimes_{\rho_{n-1}} \mathbb{R}^{n-2} \rtimes_{\rho_{n-2}} ..... \rtimes_{\rho_3} \mathbb{R}^2 \rtimes_{\rho_2} \mathbb{R}^1$ as*

$$\begin{aligned} & \mathcal{F}^n\mathcal{F}^{n-1}\mathcal{F}^{n-2}....\mathcal{F}^2\mathcal{F}^1 f(\lambda^n,\ \lambda^{n-1}, \lambda^{n-2}, ..., \lambda^2, \lambda^1) \\ = & \int_N f(X^n, X^{n-1}, ..., X^2, X^1) e^{-\,i\langle\, (\lambda^n,\, \lambda^{n-1}),(X^n,X^{n-1}),..,(\lambda^2,\, \lambda^1),(X^2,X^1)\rangle} \\ & dX^n dX^{n-1}...dX^2dX^1 \end{aligned} \quad (6)$$

for any $f \in L^1(N)$, where $X = (X^n, X^{n-1}, .., X^n, X^{n-1})$, $\mathcal{F}^d = \mathcal{F}^n\mathcal{F}^{n-1}\mathcal{F}^{n-2}....\mathcal{F}^2\mathcal{F}^1$ is the classical Fourier transform on N, $dX = dX^ndX^{n-1}...dX^2dX^1$, $\lambda = (\lambda^n,\, \lambda^{n-1}, \lambda^{n-2}, .., \lambda^2, \lambda^1)$, and

$$\langle(\lambda^n, \lambda^{n-1}), (X^n, X^{n-1}), .., (\lambda^2, \lambda^1), (X^2, X^1)\rangle = \sum_{i=1}^{n} X_i^n\lambda_i^n + \sum_{j=1}^{n-1} X_j^{n-1}\lambda_j^{n-1} +$$

$$.. + \sum_{i=1}^{2} X_i^2\lambda_i^2 + X^1\lambda^1$$

Plancherels theorem 2.1. *For any function* $f \in L^1(N)$, *we have*

$$\int_N \left|\mathcal{F}^d f(\xi^n, X^{n-1}, X^{n-2}, .., X^2, X^1)\right|^2 dX dX^{n-1} dX^{n-2} .. dX^2 dX^1$$
$$\int_N \left|\mathcal{F}^d f(\xi^n, \lambda^{n-1}, \lambda^{n-2}, .., \lambda^2, \lambda^1)\right|^2 d\xi^n d\lambda^{n-1} d\lambda^{n-2} .. d\lambda^2 d\lambda^1 \tag{7}$$

Proof: To prove this theorem, we refer to [12].

3 Notation and Results for the Solvable Lie Group AN.

3.1. Let $G = SL(n, \mathbb{R})$ be the real semi-simple Lie group and let $G = KAN$ be the Iwasawa decomposition of G, where $K = SO(n, \mathbb{R})$,and

$$N = \begin{pmatrix} 1 & * & . & . & * \\ 0 & 1 & * & . & * \\ . & . & . & . & * \\ . & . & . & . & . \\ 0 & 0 & . & 0 & 1 \end{pmatrix}, \tag{8}$$

$$A = \begin{pmatrix} a_1 & 0 & 0 & . & 0 \\ 0 & a_2 & 0 & . & 0 \\ . & . & . & . & . \\ . & . & . & . & . \\ 0 & 0 & . & 0 & a_n \end{pmatrix} \tag{9}$$

where $a_1.a_2....a_n = 1$ and $a_i \in \mathbb{R}_+^*$. The product AN is a closed subgroup of G and is isomorphic (algebraically and topologically) to the semi-direct product of A and N with N normal in AN.

Then the group AN is nothing but the group $S = N \rtimes {}_\tau A$. So the product of two elements X and$Y$ by

$$(x,\ a)(m,\ b) = (x.\tau(a)y,\ a.b) \tag{10}$$

for any $X = (x, a_1, a_2, .., a_n) \in S$ and $Y = (m, b_1, b_2, .., b_n) \in S$. Let $dnda = dmda_1da_2..da_{n-1}$ be the right haar measure on S and let $L^2(S)$ be the Hilbert space of the group S. Let $L^1(S)$ be the Banach algebra that consists of all complex valued functions on the group S, which are integrable with respect to the Haar measure of S and multiplication is defined by convolution on S as

$$g * f = \int_S f((m,b)^{-1}(n,a))g(m,b)dmdb \tag{11}$$

where $dmdb = dmdb_1db_2..db_{n-1}$ is the right Haar measure on $S = N \rtimes {}_\rho A$. Let $\Lambda = N \times A \times A$ be the group with law

$$(x,b,a)(y,c,\ d) = (x.\tau(a)y, bc, ad) \tag{12}$$

for any $(x, b, a) \in \Lambda$, and $(y, c, d) \in \Lambda$

Definition 3.1. *For every function* f defined on S, *one can define a function* $\widetilde{f}$ on Λ *as follows:*

$$\widetilde{f}(n,a,b) = f(\rho(a)n, ab) \tag{13}$$

for all $(n, a, b) \in \Lambda$. *So every function* $\psi(n, a)$ *on* S *extends uniquely as an invariant function* $\widetilde{\psi}(n,\ b,\ a)$ *on* Λ.

Remark 3.1. *The function* $\widetilde{f}$ *is invariant in the following sense:*

$$\widetilde{f}(\rho(s)n, as^{-1}, bs) = \widetilde{f}(\rho(s)n, s^{-1}a, bs) = \widetilde{f}(n,a,b) \tag{14}$$

for any $(n, a, b) \in \Lambda$ *and* $s \in A$

Lemma 3.1. *For every function* $f \in L^1(S)$ *and for every* $g \in L^1(S)$, *we have*

$$g * \widetilde{f}(n,a,b) = g *_c \widetilde{f}(n,a,b) \tag{15}$$

$$\int_{\mathbb{R}^{n-1}} \mathcal{F}_A^{n-1}\mathcal{F}^d(g * \widetilde{f})(\lambda,\mu,\nu)d\nu = \mathcal{F}_A^{n-1}\mathcal{F}^d\widetilde{f}(\lambda,\mu,0)\mathcal{F}_A^{n-1}\mathcal{F}^d g(\lambda,\mu) \tag{16}$$

for every $(n, a, b) \in \Lambda$, *where* $*$ *signifies the convolution product on* S *with respect the variables* (n, b) *and* $*_c$*signifies the commutative convolution product on the commutative group* $B = N \times A$ *with respect the variables* (n, a), *where* $\mathcal{F}_A^{n-1}$ is the Fourier transform on A.

Plancherel theorem for S. **3.1.** *For any function* $\Psi \in L^1(S)$, *we have*

$$\int\limits_{\mathbb{R}^d}\int\limits_{\mathbb{R}^{n-1}} \left|\mathcal{F}^d\mathcal{F}_A^{n-1}\Psi(\lambda,\mu)\right|^2 d\lambda d\mu = \int\limits_{AN} |\Psi(X,a)|^2 \ dXda \tag{17}$$

Go back to [12], for the prove of this theorem

4 Fourier Transform and Plancherel Theorem on $GL(n,\mathbb{R})$.

4.1. In the following we use the Iwasawa decomposition of $G = SL(n,\mathbb{R})$, to define the Fourier transform and to get Plancherel theorem on $G = SL(n,\mathbb{R})$. We denote by $L^1(G)$ the Banach algebra that consists of all complex valued functions on the group G, which are integrable with respect to the Haar measure of G and multiplication is defined by convolution on G ,

$$\phi * f(g) = \int\limits_G f(h^{-1}g)\phi(g)dg \tag{18}$$

Let $G = SL(n,\mathbb{R}) = KNA$ be the Iawsawa decomposition of G.The Haar measure dg on G can be calculated from the Haar measures dn; da and dk on N; A and K; respectively, by the formula

$$\int\limits_G f(g)dg = \int\limits_A\int\limits_N\int\limits_K f(ank)dadndk \tag{19}$$

Keeping in mind that $a^{-2\rho}$ is the modulus of the automorphism $n \to ana^{-1}$ of N we get also the following representation of dg

$$\int\limits_G f(g)dg = \int\limits_A\int\limits_N\int\limits_K f(ank)dadndk = \int\limits_N\int\limits_A\int\limits_K f(nak)a^{-2\rho}dndadk \tag{20}$$

where $\rho = \dim N = \frac{n(n+1)}{2} = 1+2+3+....+n-2+n-1+n$. Furthermore, using the relation $\int\limits_G f(g)dg = \int\limits_G f(g^{-1})dg$, we receive

$$\int\limits_K\int\limits_A\int\limits_N f(nak)a^{-2\rho}dndadk = \int\limits_K\int\limits_A\int\limits_N f(kan)a^{2\rho}dndadk \tag{21}$$

Definition 4.1. *The Fourier transform of a function $f \in C^{\infty}(K)$ is defined as*

$$Tf(\gamma) = \int_K f(x)\gamma(x^{-1})dx \tag{22}$$

where T is the Fourier transform on K, and $\gamma \in \widehat{K}$ ($\widehat{K}$ is the set of irreducible unitary representations of K)

Theorem (A. Cerezo) 4.1. *Let $f \in C^{\infty}(K)$, then we have the inversion of the Fourier transform*

$$f(x) = \sum_{\gamma \in \widehat{K}} d\gamma tr[Tf(\gamma)\gamma(x)] \tag{23}$$

and the Plancherel formula

$$\|f(x)\|_2^2 = \int |f(x)|^2 \, dx = \sum_{\gamma \in \widehat{K}} d_\gamma \, \|Tf(\gamma)\|_{H.S}^2 \tag{24}$$

for any $f \in L^1(K)$, see $[2, P.562-563]$,where $\|Tf(\gamma)\|_{H.S}^2$ is the norm of Hilbert-Schmidt of the operator $Tf(\gamma)$.

Definition 4.2. *For any function $f \in \mathcal{D}(G)$, we can define a function $\Upsilon(f)$on $G \times K$ by*

$$\Upsilon(f)(g, k_1) = \Upsilon(f)(kna, k_1) = f(gk_1) = f(knak_1) \tag{25}$$

for $g = kna \in G$, and $k_1 \in K$. The restriction of $\Upsilon(f) * \psi(g, k_1)$ on $K(G)$ is $\Upsilon(f) * \psi(g, k_1) \downarrow_{K(G)} = f(nak_1) \in \mathcal{D}(G)$, and $\Upsilon(f)(g, k_1) \downarrow_K = f(g, I_K) = f(kna) \in \mathcal{D}(G)$

Remark 4.1. *The function $\Upsilon(f)$ is invariant in the following sense*

$$\Upsilon(f)(gh, h^{-1}k_1) = f(gk_1) = f(knak_1) \tag{26}$$

Definition 4.3. *Let f and ψ be two functions belong to $\mathcal{D}(G)$, then we can define the convolution of $\Upsilon(f)$ and ψ on $G \times K$ as*

$$\begin{aligned} \Upsilon(f) * \psi(g, k_1) &= \int_G \Upsilon(f)(gg_2^{-1}, k_1)\psi(g_2)dg_2 \\ &= \int_K \int_N \int_A \Upsilon(f)(knaa_2^{-1}n_2^{-1}k^{-1}k_1)\psi(k_2n_2a_2)dk_2dn_2da_2 \end{aligned}$$

So we get

$$\begin{aligned}\Upsilon(f)*\psi(g,k_1) \quad \downarrow \quad _{K(G)} &= \Upsilon(f)*\psi(I_K na,k_1)\\ &= \int\limits_K\int\limits_N\int\limits_A f(naa_2^{-1}n_2^{-1}k^{-1}k_1)\psi(k_2n_2a_2)dk_2dn_2da_2\end{aligned}$$

where T is the Fourier transform on K, and I_K is the identity element of K. Denote by $\mathcal{F}$ is the Fourier transform on AN

Definition 4.4. *If $f\in\mathcal{D}(G)$ and let $\Upsilon(f)$ be the associated function to f , we define the Fourier transform of $\Upsilon(f)(g,k_1)$ by*

$$\begin{aligned}&T\mathcal{F}\Upsilon(f))(I_K,\xi,\lambda,\gamma)=T\mathcal{F}\Upsilon(f))(I_K,\xi,\lambda,\gamma)\\ =\ &\int_K\int_N\int_A\sum_{\delta\in\widehat{K}}d_\delta tr[\int_K\Upsilon(f)(kna,k_1)\delta(k^{-1})dk]a^{-i\lambda}e^{-\ i\langle\ \xi,\ n\rangle}\ \gamma(k_1^{-1})dadndk_1\\ =\ &\int_N\int_A\int_K\Upsilon(f)(I_Kna,k_1)a^{-i\lambda}e^{-\ i\langle\ \xi,\ n\rangle}\ \gamma(k_1^{-1})dadndk_1\\ =\ &\int_N\int_A\int_K f(nak_1)a^{-i\lambda}e^{-\ i\langle\ \xi,\ n\rangle}\ \gamma(k_1^{-1})dadndk_1 \qquad (27)\end{aligned}$$

Theorem 4.2. (***Plancherel's Formula for the Group*** $G=SL(n,\mathbb{R})$**.** *For any function $f\in L^1(G)\cap L^2(G)$,we get*

$$\begin{aligned}\int|f(g)|^2\,dg \ &= \int_K\int_N\int_A|f(kna)|^2\,dadndk\\ &= \sum_{\gamma\in\widehat{K}}d_\gamma\int\limits_{\mathbb{R}^d}\int\limits_{\mathbb{R}^{n-1}}\|T\mathcal{F}f(\lambda,\xi,\gamma)\|^2_{H.S}\,d\lambda d\xi\\ &= \int\limits_{\mathbb{R}^d}\int\limits_{\mathbb{R}^{n-1}}\sum_{\gamma\in\widehat{K}}d_\gamma\|T\mathcal{F}f(\lambda,\xi,\gamma)\|^2_{H.S}\,d\lambda d\xi \qquad (28)\end{aligned}$$

where I_A,I_N and *I_K are the identity elements of A, N and K respectively, $\mathcal{F}$ is the Fourier transform on AN and T is the Fourier transform on K*

For the proof of this theorem see [13]

New Group. Let $GL(n,\mathbb{R})$ be the general linear group consisting of all matrices of the form

$$GL(n,\mathbb{R})=\{X=\left(\ a_{ij}\ \right),a_{ij}\in\mathbb{R},\ \ \prec 1\prec n,,\ \ \prec j\ \prec n,\ and\ \det A\neq 0\} \qquad (29)$$

As a manifold, $GL(n,\mathbb{R})$ is not connected but rather has two connected components: the matrices with positive determinant and the ones with negative determinant which is denoted by $GL_-(n,\mathbb{R})$. The identity component, denoted by $GL_+(n,\mathbb{R})$, consists of the real $n\times n$ matrices with positive determinant. This is also a Lie group of dimension n^2; it has the same Lie algebra as $GL(n,\mathbb{R})$.

The group $GL(n,\mathbb{R})$ is also noncompact. The maximal compact subgroup of $GL(n,\mathbb{R})$ is the orthogonal group $O(n)$, while the maximal compact subgroup of $GL_+(n,\mathbb{R})$ is the special orthogonal group $SO(n)$. As for $SO(n)$, the group $GL_+(n,\mathbb{R})$ is not simply connected

Theorem 4.3. *$GL_-(n,\mathbb{R})$ is group isomorphic onto $GL_+(n,\mathbb{R})$*

For the proof of this theorem see [13].

As well known the group $GL_+(n,\mathbb{R})$ is isomorphic onto the direct product of the two groups $SL(n,\mathbb{R})$ and $\mathbb{R}_+^*$, *i.e* $GL_+(n,\mathbb{R})$ =$SL(n,\mathbb{R})\times\ \mathbb{R}_+^*$ and $GL(n,\mathbb{R})$ =$GL_-(n,\mathbb{R})\ \cup GL_+(n,\mathbb{R}) = (SL(n,\mathbb{R})\times\ \mathbb{R}_+^*)\cup(SL(n,\mathbb{R})\times\ \mathbb{R}_+^*)$. Our aim result is

Plancherel theorem 4.4. *Let $\mathcal{F}_+^*$ be the Fourier transform on $GL_+(n,\mathbb{R}$ the we get*

$$\begin{aligned}\int_{GL_+(n,\mathbb{R})}|f(g,t)|^2\,dg\frac{dt}{t} &= \int\limits_{\mathbb{R}_+^*}\int_K\int_N\int_A|f(kna,t)|^2\,dadndk\frac{dt}{t}\\ &= \sum_{\gamma\in\widehat{K}}d_\gamma\int\limits_{\mathbb{R}}\int\limits_{\mathbb{R}^d}\int\limits_{\mathbb{R}^{n-1}}\left\|\mathcal{F}_+^*T\mathcal{F}f(\lambda,\xi,\gamma,\eta)\right\|_{H.S}^2d\lambda d\xi d\eta\\ &= \int\limits_{\mathbb{R}}\int\limits_{\mathbb{R}^d}\int\limits_{\mathbb{R}^{n-1}}\sum_{\gamma\in\widehat{K}}d_\gamma\left\|\mathcal{F}_+^*T\mathcal{F}f(\lambda,\xi,\gamma,\eta)\right\|_{H.S}^2d\lambda d\xi d\eta\end{aligned}\tag{30}$$

for any $f\in L^1(GL_+(n,\mathbb{R})\cap L^2(GL_+(n,\mathbb{R})$. For the proof of this theorem see [13].

Corollary 4.1. *Let* f be a function belongs $L^1(GL(n,\mathbb{R}) \cap L^2(GL(n,\mathbb{R})$

$$\begin{aligned}
&\int_{GL(n,\mathbb{R})} |f(g,t)|^2 \, dg \frac{dt}{t} \\
= &\int_{GL_-(n,\mathbb{R}) \cup GL_+(n,\mathbb{R})} |f(g,t)|^2 \, dg \frac{dt}{t} = 2 \int_{GL_+(n,\mathbb{R})} |f(g,t)|^2 \, dg \frac{dt}{t} \\
= &\; 2 \int\limits_{\mathbb{R}_+^*} \int_K \int_N \int_A |f(kna,t)|^2 \, da dn dk \frac{dt}{t} \\
= &\; 2 \sum_{\gamma \in \widehat{K}} d_\gamma \int\limits_{\mathbb{R}} \int\limits_{\mathbb{R}^d} \int\limits_{\mathbb{R}^{n-1}} \left\| \mathcal{F}_+^* T \mathcal{F} f(\lambda, \xi, \gamma, \eta) \right\|_{H.S}^2 d\lambda d\xi d\eta \\
= &\; 2 \int\limits_{\mathbb{R}} \int\limits_{\mathbb{R}^d} \int\limits_{\mathbb{R}^{n-1}} \sum_{\gamma \in \widehat{K}} d_\gamma \left\| \mathcal{F}_+^* T \mathcal{F} f(\lambda, \xi, \gamma, \eta) \right\|_{H.S}^2 d\lambda d\xi d\eta \qquad (31)
\end{aligned}$$

Remark 2.1. this corollary explains the Fourier transform and Plancherel formula on the non connected Lie group $GL(n,\mathbb{R})$.

5 Fourier Transform and Plancherel Theorem for $GA(n,\mathbb{R})$

5.1. We begin by the group $GA_+(n,\mathbb{R}) = \mathbb{R}^n \rtimes_\rho GL_+(n,\mathbb{R})$ to define the Fourier transform and to prove the Plancherel formula on the group $GA(n,\mathbb{R})$,where ρ is the group homomorphism from $GL_+(n,\mathbb{R})$ into the group $Aut(\mathbb{R}^n)$ of all automorphisms of the real vector group $\mathbb{R}^n$. Let $H_+ = \mathbb{R}^n \times GL(n,\mathbb{R})_+ \times GL_+(n,\mathbb{R})$ be the Lie group with the following law

$$(A,X,Y)(B,P,Q) = (A + \rho(Y)B, XB, YQ) \qquad (32)$$

for all $(A,X,Y) \in H_+$, $(C,D,Y) \in H_+$. Denote by $L_+ = \mathbb{R}^n \times GL_+(n,\mathbb{R})$ the lie group which is direct product of the two groups $\mathbb{R}^n$ and $GL_+(n,\mathbb{R})$. In this case the group $GA_+(n,\mathbb{R})$ can be identified with the closed subgroup $\mathbb{R}^n \times \{0\} \times_\rho GL_+(n,\mathbb{R})$ of H_+ and L_+ with the subgroup $\mathbb{R}^n \times GL_+(n,\mathbb{R}) \times \{0\}$ of H_+

Definition 5.1. *For every function* f defined on $GA_+(n,\mathbb{R})$, *one can define a function* $\widetilde{f}$ on H_+ *as follows:*

$$\widetilde{f}(A,X,Y) = f(\rho(X)A, XY) \qquad (33)$$

for all $(A,X,Y)\in H_+$, *where* $X=(k_1n_1a_1,t_1)$, $Y=(k_2n_2a_2,t_2)$, $(k_1,k_2)\in K\times K$, $(n_1,n_2)\in N\times N$, *and* $(t_1,t_2)\in\mathbb{R}_+^*\times\mathbb{R}_+^*$. *So every function* $\psi(A,Y)\in GA_+(n,\mathbb{R})$ *extends uniquely as an invariant function* $\widetilde{\psi}(A,X,Y)$ *on* H_+.

Remark 5.1. *The function* $\widetilde{f}$ *is invariant in the following sense:*

$$\widetilde{f}(\rho(T)A,XT^{-1},TY)=\widetilde{f}(A,X,Y) \tag{34}$$

for any $(A,X,Y)\in H_+$ *and* $T\in GL_+(n,\mathbb{R})$

Lemma 5.1. *For every function* $f\in L^1(GA_+(n,\mathbb{R}))$ *and for every* $g\in L^1(GA_+(n,\mathbb{R})$, *we have*

$$g*\widetilde{f}(A,X,Y)=\widetilde{f}*_c g(A,X,Y) \tag{35}$$

for every $(A,X,Y)\in H_+$, *where* $*$ *signifies the convolution product on* $GA_+(n,\mathbb{R})$*with respect the variables* (A,Y) *and* $*_c$*signifies the convolution product on* $L_+=\mathbb{R}^n\times GL_+(n,\mathbb{R})$ *with respect the variables* (A,X),

Proof: In fact for any f and g belong $L^1(GA_+(n,\mathbb{R}))$ we have

$$\begin{aligned}
g*\widetilde{f}(A,X,Y) &= \int\limits_{\mathbb{R}^n}\int\limits_{GL_+(n,\mathbb{R})}\widetilde{f}((B,Q)^{-1}(A,X,Y))g(B,Q)dBdQ\\
&= \int\limits_{\mathbb{R}^n}\int\limits_{GL_+(n,\mathbb{R})}\widetilde{f}\left[(\rho(Q^{-1})(-B),Q^{-1})(A,X,Y)\right]g(B,Q)dBdQ\\
&= \int\limits_{\mathbb{R}^n}\int\limits_{GL_+(n,\mathbb{R})}\widetilde{f}\left[\rho(Q^{-1})(A-B),X,Q^{-1}Y\right]g(B,Q)dBdQ\\
&= \int\limits_{\mathbb{R}^n}\int\limits_{GL_+(n,\mathbb{R})}\widetilde{f}\left[A-B,XQ^{-1},Y\right]g(B,Q)dBdQ\\
&= \widetilde{f}*_c g(A,X,Y)
\end{aligned} \tag{36}$$

Definition 5.1. *If* $f\in L^1(GA_+(n,\mathbb{R}))$, *one can define the Fourier transform of* f *as*

$$\begin{aligned}
&\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}f(\mu,(\gamma,\xi,\lambda,\eta))\\
&= \int\limits_{\mathbb{R}^n}\int_{\mathbb{R}_+^*}\int_N\int_A\int_K f(A,ank,t)e^{-i\langle\,\mu,\,A\rangle}t^{-i\eta}a^{-i\lambda}e^{-i\langle\,\xi,\,n\rangle}\,\gamma(k^{-1})dAdadndk\frac{dt}{t}
\end{aligned}$$

here $\mathcal{F}_{GL_+}$ denotes the Fourier transform on $GL_+(n,\mathbb{R})$ and Let $\mathcal{F}_{\mathbb{R}^n}$ be the Fourier transform on $\mathbb{R}^n$

Corollary 5.1. *For any function $f \in L^1(GA_+(n,\mathbb{R}))$, we have*

$$\begin{aligned}
&\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}}\sum_{\delta\in\widehat{K}} d_\delta tr[\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}(g*\widetilde{f})(\mu,(\gamma,\xi,\lambda,,\eta),(\delta,\nu,\sigma,\omega)]d\nu d\sigma d\omega \\
&= [\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}(\widetilde{f})(\mu,(\gamma,\xi,\lambda,,\eta),I_{GL_+})\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}(g)(\mu,(\gamma,\xi,\lambda,\eta))] \qquad (37)
\end{aligned}$$

The proof of this theorem results immediately from lemma **4.1**

Plancherel Theorem 5.1. *For any function $f \in L^1(GA_+(n,\mathbb{R}))$, we have*

$$\begin{aligned}
&= \int_{\mathbb{R}^n}\int_{GL_+(n,\mathbb{R})} |f(B,Q)|^2\, dBdQ \\
&= \int_{GA_+(n,\mathbb{R})} |f(A,g,t)|^2\, dBdg\frac{dt}{t} \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}_+^*}\int_K\int_N\int_A |f(A,kna,t)|^2\, dBdadndk\frac{dt}{t} \\
&= \sum_{\gamma\in\widehat{K}} d_\gamma \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}} \left\|\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}f(\mu,(\gamma,\xi,\lambda,\eta))\right\|_{H.S}^2 d\mu d\lambda d\xi d\eta \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}}\sum_{\gamma\in\widehat{K}} d_\gamma \left\|\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}f(\mu,(\gamma,\xi,\lambda,\eta)\right\|_{H.S}^2 d\mu d\lambda d\xi d\eta \qquad (38)
\end{aligned}$$

Proof: Let $\widetilde{\overset{\vee}{f}}$ the function defined as

$$\widetilde{\overset{\vee}{f}}(A,X,Y) = \overset{\vee}{f}(\rho(X)A,XY) = \overline{f[(\rho(X)A,XY)^{-1}]} \qquad (39)$$

Then we have

$$
\begin{aligned}
& f * \overset{\widetilde{\vee}}{f}(0, I_{GL_+}, I_{GL_+}) \\
= & \int_{GA_+(n,\mathbb{R})} \overset{\widetilde{\vee}}{f}[(B,Q)^{-1}(0, I_{GL_+}, I_{GL_+})] f(B,Q) dB dQ \\
= & \int_{\mathbb{R}^n} \int_{GL_+(n,\mathbb{R})} \overset{\widetilde{\vee}}{f}\left[\rho(Q^{-1})(-B), I_{GL_+}, Q^{-1} I_{GL_+}\right] f(B,Q) dB dQ \\
= & \int_{\mathbb{R}^n} \int_{GL_+(n,\mathbb{R})} \overset{\widetilde{\vee}}{f}\left[-B, Q^{-1}, I_{GL_+}\right] f(B,Q) dB dQ \\
= & \int_{\mathbb{R}^n} \int_{GL_+(n,\mathbb{R})} \overset{\vee}{f}\left[\rho(Q^{-1})(-B), Q^{-1}\right] f(B,Q) dB dQ \\
= & \int_{\mathbb{R}^n} \int_{GL_+(n,\mathbb{R})} \overline{f(B,Q) f}(B,Q) dB dQ \\
= & \int_{\mathbb{R}^n} \int_{GL_+(n,\mathbb{R})} |f(B,Q)|^2 \, dB dQ \\
= & \int_{\mathbb{R}^n} \int_{GA_+(n,\mathbb{R})} |f(B,g,t)|^2 \, dB dg \frac{dt}{t} \\
= & \int_{\mathbb{R}^n} \int_{GA_+(n,\mathbb{R})} |f(B,ank,t)|^2 \, dB dn da dk \frac{dt}{t} \qquad (40) \\
= & \int_{\mathbb{R}^n} \int_{GA_+(n,\mathbb{R})} |f(B,kna,t)|^2 \, dB dn da dk \frac{dt}{t} \qquad (41)
\end{aligned}
$$

where I_{GL_+} is the identity element of $GL_+(n,\mathbb{R})$. In other hand, we get

$$
\begin{aligned}
& f * \widetilde{\vee f}(0, I_{GL_+}, I_{GL_+}) \\
= & \int_{\mathbb{R}^n} \int_{\mathbb{R}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^{n-1}} \sum_{\gamma \in \widehat{K}} d_\gamma \sum_{\delta \in \widehat{K}} d_\delta tr[\mathcal{F}_{\mathbb{R}^n} \mathcal{F}_{GL_+} (f * \widetilde{\vee f})(\mu, (\gamma, \xi, \lambda, , \eta), (\delta, \nu, \sigma, \omega)] \\
& d\mu d\lambda d\xi d\eta d\nu d\sigma d\omega \\
= & \int_{\mathbb{R}^n} \int_{\mathbb{R}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^{n-1}} \sum_{\gamma \in \widehat{K}} d_\gamma tr[\mathcal{F}_{\mathbb{R}^n} \mathcal{F}_{GL_+} (\widetilde{\vee f})(\mu, (\gamma, \xi, \lambda, , \eta), I_{GL_+})] \mathcal{F}_{\mathbb{R}^n} \mathcal{F}_{GL_+} f(\mu, \lambda, \xi, \gamma, \eta) \\
& d\mu d\lambda d\xi d\eta
\end{aligned}
$$

Calculate the formula $\mathcal{F}_{\mathbb{R}^n} \mathcal{F}_{GL_+} (\widetilde{\vee f})(\mu, (\gamma, \xi, \lambda, \eta), I_{GL_+})$

$$
\begin{aligned}
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}} \mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}(\widetilde{\overset{\vee}{f}})(\mu,(\mu,(\gamma,\xi,\lambda,,\eta),I_{GL_+}) \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}}\int_{\mathbb{R}^n}\int_{\mathbb{R}_+^*}\int_N\int_A\int_K \widetilde{\overset{\vee}{f}}(A,(kna,t),I_{GL_+}) \\
&\quad e^{-\,i\langle\,\mu,\,A\rangle}t^{-i\eta}a^{-i\lambda}e^{-\,i\langle\,\xi,\,n\rangle}\,\gamma(k^{-1})dAdkdnda\frac{dt}{t}d\mu d\lambda d\xi d\eta \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}}\int_{\mathbb{R}^n}\int_{\mathbb{R}_+^*}\int_N\int_A\int_K \overset{\vee}{f}(\rho(ank,t)A,(ank.t)) \\
&\quad e^{-\,i\langle\,\mu,\,A\rangle}t^{-i\eta}a^{-i\lambda}e^{-\,i\langle\,\xi,\,n\rangle}\,\gamma(k^{-1})dkdAdadn\frac{dt}{t}d\mu d\lambda d\xi d\eta \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}}\int_{\mathbb{R}^n}\int_{\mathbb{R}_+^*}\int_N\int_A\int_K \overline{f((\rho(ank,t)A,(ank.t))^{-1})} \\
&\quad e^{-\,i\langle\,\mu,\,A\rangle}t^{-i\eta}a^{-i\lambda}e^{-\,i\langle\,\xi,\,n\rangle}\,\gamma(k^{-1})dkdAdadn\frac{dt}{t}d\mu d\lambda d\xi d\eta \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}}\int_{\mathbb{R}^n}\int_{\mathbb{R}_+^*}\int_N\int_A\int_K \overline{f((-A,(k^{-1}n^{-1}a^{-1},t^{-1}))} \\
&\quad e^{-\,i\langle\,\mu,\,A\rangle}t^{-i\eta}a^{-i\lambda}e^{-\,i\langle\,\xi,\,n\rangle}\,\gamma(k^{-1})dk]dAdadn\frac{dt}{t}d\mu d\lambda d\xi d\eta \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}}\int_{\mathbb{R}^n}\int_{\mathbb{R}_+^*}\int_N\int_A\int_K \overline{f((-A,(kna,t))} \\
&\quad \overline{e^{-\,i\langle\,\mu,\,A\rangle}t^{-i\eta}a^{-i\lambda}e^{-\,i\langle\,\xi,\,n\rangle}}\,\gamma^*(k^{-1})dkdAdadn\frac{dt}{t}d\mu d\lambda d\xi d\eta \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}} \overline{\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}(f)(\mu,(\gamma^*,\xi,\lambda,\eta))}d\mu d\lambda d\xi d\eta
\end{aligned}
$$

Finally, we get

$$
\begin{aligned}
& f * \overset{\widetilde{\vee}}{f}(0, I_{GL_+}, I_{GL_+}) \\
= & \int\limits_{\mathbb{R}^n} \int\limits_{GL_+(n,\mathbb{R})} |f(A,Q)|^2 \, dAdQ \\
= & \int\limits_{\mathbb{R}^n} \int_{GA_+(n,\mathbb{R})} |f(A,g,t)|^2 \, dAdg\frac{dt}{t} \\
= & \int\limits_{\mathbb{R}^n} \int_{GA_+(n,\mathbb{R})} |f(A,ank,t)|^2 \, dAdndadk\frac{dt}{t} \\
= & \int\limits_{\mathbb{R}^n} \int_{GA_+(n,\mathbb{R})} |f(A,kna,t)|^2 \, dAdndadk\frac{dt}{t} \qquad (42) \\
= & \int\limits_{\mathbb{R}^n} \int\limits_{\mathbb{R}} \int\limits_{\mathbb{R}^d} \int\limits_{\mathbb{R}^{n-1}} \sum_{\gamma \in \widehat{K}} d_\gamma tr[\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}(\overset{\widetilde{\vee}}{f})(\mu,(\gamma,\xi,\lambda,\eta),I_{GL_+})\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}f(\mu,\gamma,\xi,\lambda,\eta)] \\
& d\mu d\lambda d\xi d\eta \\
= & \int\limits_{\mathbb{R}^n} \int\limits_{\mathbb{R}} \int\limits_{\mathbb{R}^d} \int\limits_{\mathbb{R}^{n-1}} \sum_{\gamma \in \widehat{K}} d_\delta tr[\overline{\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}(f)(\mu,(\gamma^*,\xi,\lambda,\eta))}\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}f(\mu,(\gamma,\xi,\lambda,,\eta))] \\
= & \int\limits_{\mathbb{R}^n} \int\limits_{\mathbb{R}} \int\limits_{\mathbb{R}^d} \int\limits_{\mathbb{R}^{n-1}} \sum_{\gamma \in \widehat{K}} d_\delta \left\| \mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+}f(\mu,(\gamma,\xi,\lambda,,\eta) \right\|_{H.S}^2 d\mu d\lambda d\xi d\eta \qquad (43)
\end{aligned}
$$

Hence the proof of our theorem. Now we can state our final result

Theorem 5.2. *For any function* $f \in L^1(GA(n,\mathbb{R}))$, *we get*

$$
\begin{aligned}
&= \int_{\mathbb{R}^n}\int_{GL(n,\mathbb{R})} |f(A,Q)|^2\, dAdQ \\
&= 2\int_{\mathbb{R}^n}\int_{GL_+(n,\mathbb{R})} |f(A,Q)|^2\, dAdQ \qquad (44)\\
&= 2\int_{GA_+(n,\mathbb{R})} |f(A,g,t)|^2\, dAdg\frac{dt}{t} \\
&= 2\int_{\mathbb{R}^n}\int_{\mathbb{R}^*_+}\int_K\int_N\int_A |f(A,kna,t)|^2\, dAdadndk\frac{dt}{t} \\
&= 2\sum_{\gamma\in\widehat{K}} d_\gamma \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}} \left\|\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+} f(\mu,(\gamma,\xi,\lambda,,\eta))\right\|^2_{H.S} d\mu d\lambda d\xi d\eta \\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}}\int_{\mathbb{R}^d}\int_{\mathbb{R}^{n-1}} 2\sum_{\gamma\in\widehat{K}} d_\gamma \left\|\mathcal{F}_{\mathbb{R}^n}\mathcal{F}_{GL_+} f(\mu,(\gamma,\xi,\lambda,,\eta)\right\|^2_{H.S} d\mu d\lambda d\xi d\eta \qquad (45)
\end{aligned}
$$

Conclusion. I believe that the results of this paper will be a guideline to study the Fourier analysis on non connected locally compact Lie groups.

References

[1] M. F. Atiyah; (1970), *Resolution of Singularities and Division of Distributions*, Comm.on Pure and App. Math, Vol, 23, 145-150.

[2] V. Bargmann; (1947), *Irreducible Unitary Representations of the Lorentz Group*, Ann. of Math. 48, 568-640.

[3] A. Cerezo and F. Rouviere, (1969)*"Solution Elemetaire d'un Operator Differentielle Lineare Invariant Agauch sur un Group de Lie Reel Compact"* Annales Scientiques de E.N.S. 4 serie, tome 2, n^{o}4,p 561-581.

[4] K. El- Hussein., (2009), *A Fundamental Solution of an Invariant Differential Operator on the Heisenberg Group*, International Mathematical Forum, 4, no. 12, 601- 612.

[5] K. El- Hussein., (2011), *On the Left Ideals of Group Algebra of thje Affine Group.* International Mathematical Forum, Vol 6. no. 4, 193-204.

[6] K. El- Hussein., (1989), *Opérateurs Différéntiels Invariants sur les Groupes de Deplacements,* Bull. Sc. Math. 2^e series 113, 89-117.

[7] K. El- Hussein., (2009), *Eigendistributions for the Invariant Differential Operators on the Affine Group*, Int. Journal of Math. Analysis, Vol. 3, no. 9 , 419-429.

[8] K. El- Hussein, (2010), *Note on the Solvability of Lewy Operator*, Int. Math. Forum, 5, no. 8, 389 - 393.

[9] K. El- Hussein., *(*2010), *Fourier transform and invariant differential operators on the solvable Lie group G4*, in Int. J. Contemp. Maths Sci. 5. No. 5-8, 403-417.

[10] K. El- Hussein., *(*2013), *Abstract Harmonic Analysis and Ideals of Banach Algebra on 3-Step Nilpotent Lie Group*, International Journal of Engineering Research & Technology (IJERT), Vol. 2 Issue 11, November.

[11] K. El- Hussein., (2014), *Abstract Harmonic Analysis on Spacetime*, arXiv:1404.1600v1 [math-ph] 6 Apr.

[12] K. El- Hussein., (2014), *Abstract Harmonic Analysis and Group Algebra on the Group AN*, arXiv:1404.3361v1 [math.GR] 13 Apr.

[13] K. El- Hussein., (2014), *Abstract Harmonic Analysis on the General Linear Group* $GL(n, \mathbb{R})$, arXiv:1404.5535v1 [math.RT] 20 Apr.

[14] Harish-Chandra; (1952), *Plancherel Formula for* 2×2 *Real Unimodular Group*, Proc. nat. Acad. Sci. U.S.A., vol. 38, pp. 337-342.

[15] Harish-Chandra; (1954), *The Plancherel Formula for Complex Semisimple Lie Group*, Trans. Amer. Mth. Soc., vol. 76, pp. 485- 528.

[16] Harish-Chandra; (1957), *A Formula for Semisimple Lie Groups*, Amer. J. Math. 79, 733-760.

[17] S. Helgason., (1984), *Groups and Geometric Analysis*, Academic Press.

[18] S. Helgason., (2005), *The Abel, Fourier and Radon Transforms on Symmetric Spaces*. Indagationes Mathematicae. 16, 531-551.

[19] L. Hormander, (1983), *The analysis of Linear Partial Differential Operator I*", Springer-Verlag.

[20] D. Oldfeld and A. Kontorovich; (2011), *On the Determination of the Plancherel Measure for Lebedev-Whittaker transforms on* $GL(n, \mathbb{R})$ arXiv:1102.5086v1 [math.NT] 24 Feb.

[21] L. Pukanszky; (1964), *The Plancherel Formula for the Universal Covering Group of* $SL(\mathbb{R}, 2)$, Math. Ann. 156, 96-143.

[22] Roberto Camporesi*;* (1979), *The Helgason Fourier Transform for Homogeneous Vector Bundles Over Riemannian Symmetric spaces,* pacific journal of mathematics, Vol. 179, No. 2.

[23] F. Treves., (1962), *Linear Partial Differential Equations with Constant Coefficients,* Gordon and Breach.

[24] W.Rudin., (1962), *Fourier Analysis on Groups,* Interscience Publishers, New York, NY.G.

[25] Warner., (1970), *Harmonic Analysis on Semi-Simple Lie Groups*, Springer-verlag Berlin heidel;berg New york.

[26] N.R. Wallach; (1973), *Harmonic Analysis on Homogeneous Spaces*, Marcel Dekker, INC, New York.